

Reinforce Infection Control In Light of Coronavirus

The “COVID-19” coronavirus outbreak will put the spotlight on ways to limit the spread of respiratory infections.

This coronavirus is thought to be transmitted by respiratory secretions...similar to other coronaviruses, such as SARS and MERS.

Put the current risk in perspective to help calm worried patients...since it doesn't yet come close to the overall impact of seasonal influenza.

Over 35 MILLION patients got the flu last season...with about 500,000 hospitalizations and 34,000 deaths.

Emphasize getting a flu vaccine each year...and use this coronavirus outbreak as an opportunity to stress infection control.

Reinforce proper hand hygiene. Explain that frequent handwashing with soap and water is the best bet. Advise washing for 20 seconds...about the time it takes to sing the “Happy Birthday” song twice.

If that's not practical, suggest using a hand sanitizer with at least 60% alcohol. They're effective against enveloped viruses, such as coronaviruses...but aren't yet proven to kill COVID-19.

Use a disinfectant spray or wipe on exam tables, phones, etc.

Clear up mask confusion. Caution that unnecessary use is leading to shortages. Plus masks only work along with hand hygiene.

Discourage masks for most people withOUT symptoms. Masks are too loose to keep all germs out...and viruses can get in through the eyes.

If patients ask about N95 respirators, point out these masks require fitting...and are hard to wear for more than a few minutes.

But it's okay if patients WITH respiratory symptoms want to wear a surgical mask to help contain secretions...especially if in contact with infants, elderly, or immunocompromised people.

And ask patients WITH respiratory symptoms to call before coming in...if they recently traveled to an outbreak area or had contact with a COVID-19 patient.

Expect the guidance for patients with suspected or confirmed COVID-19 to evolve as we learn more.

4/14/20 – This article was published before CDC released recommendations about cloth face coverings in public settings, available here. Please keep in mind that guidance is changing quickly. Refer to the CDC for the most current recommendations.

3/13/20 - This article was revised to reflect the current population risk in North America. Please keep in mind that this is a rapidly changing landscape.

Key References:

- www.who.int/emergencies/diseases/novel-coronavirus-2019 (2-26-20)
- www.cdc.gov/coronavirus/2019-ncov/index.html (2-26-20)
- www.cdc.gov/coronavirus/2019-ncov/hcp/healthcare-supply-ppe.html (2-26-20)
- www.who.int/emergencies/diseases/novel-coronavirus-2019/advice-for-public/when-and-how-to-use-masks (2-26-20)

Cite this document as follows: Article, Reinforce Infection Control In Light of Coronavirus, Prescriber Insights: APRN, March 2020

The content of this article is provided for educational and informational purposes only, and is not a substitute for the advice, opinion or diagnosis of a trained medical professional. If your organization is interested in an enterprise subscription, email sales@trchealthcare.com.

© 2024 Therapeutic Research Center (TRC). TRC and Prescriber Insights: APRN and the associated logo(s) are trademarks of Therapeutic Research Center. All Rights Reserved.

Prescriber's Letter. March 2020, No. 360301

Cite this document as follows: Article, Reinforce Infection Control In Light of Coronavirus, Prescriber Insights: APRN, March 2020

The content of this article is provided for educational and informational purposes only, and is not a substitute for the advice, opinion or diagnosis of a trained medical professional. If your organization is interested in an enterprise subscription, email sales@trhealthcare.com.

© 2024 Therapeutic Research Center (TRC). TRC and Prescriber Insights: APRN and the associated logo(s) are trademarks of Therapeutic Research Center. All Rights Reserved.